

# BMJ Open

## A Qualitative, Grounded Theory Exploration of Patients' Experience of Early Mobilisation, Rehabilitation and Recovery after Critical Illness.

Journal:	<i>BMJ Open</i>
Manuscript ID	bmjopen-2018-026348.R2
Article Type:	Research
Date Submitted by the Author:	07-Jan-2019
Complete List of Authors:	Corner, Evelyn; Brunel University Department of Clinical Sciences; The Royal Brompton and Harefield NHS Trust Murray, Eleanor; University of Oxford Said Business School, Said Business School Brett, Stephen; Imperial College London Department of Surgery and Cancer; Imperial College Healthcare NHS Trust, Anaesthetics and Intensive Care
<b>Primary Subject Heading</b>:	Intensive care
Secondary Subject Heading:	Rehabilitation medicine, Qualitative research
Keywords:	INTENSIVE & CRITICAL CARE, REHABILITATION MEDICINE, PATIENT EXPERIENCE, RECOVERY, PHYSIOTHERAPY

SCHOLARONE™  
Manuscripts

**Title:** A Qualitative, Grounded Theory Exploration of Patients' Experience of Early Mobilisation, Rehabilitation and Recovery after Critical Illness.

**Authors:** Dr Evelyn J. Corner<sup>1,2,3</sup>, Dr Eleanor J. Murray<sup>4</sup>, Professor Stephen J. Brett<sup>5,6</sup>

1. Department of Clinical Sciences, Brunel University London, UK. Phone number: +44 1895 268489
2. Chelsea and Westminster NHS Foundation Trust, London, UK
3. The Royal Brompton and Harefield NHS Foundation Trust, London, UK.
4. Saïd Business School, University of Oxford, Oxford, UK. Phone number: +44 (0)1865 614828
5. Department of Surgery and Cancer, Imperial College London, UK
6. Centre for Perioperative Medicine and Critical Care Research, Imperial College NHS Healthcare Trust, Hammersmith Hospital, London, UK. Phone number: +44 208 383 4521

**Corresponding author:**

Email: [evelyn.corner@brunel.ac.uk](mailto:evelyn.corner@brunel.ac.uk)

Phone number: +44 1895 268489

**Financial support:** This study was kindly funded by The Joint Research Committee at Chelsea and Westminster Hospital on behalf of a joint collaboration between the Chelsea and Westminster Medical School Research Trust (Registered Charity No. 247808) and Chelsea and Westminster Health Charity, and Brunel University London. EJM acknowledges the support of The Health Foundation. SJB is grateful for the support of the National Institute for Health Research (NIHR) comprehensive Biomedical Research Centre based at Imperial College Healthcare NHS Trust and Imperial College London. The views expressed are those

1  
2  
3 of the authors and not necessarily those of the NHS, The Health Foundation, the NIHR or the  
4  
5 Department of Health.  
6  
7

8  
9 **Competing interests:** No competing interests to declare.  
10

11  
12 **Authors contribution to the study:** EJC, SJB and EJM were involved in the methodological  
13  
14 design, data analysis and writing and reviewing the manuscript. EJC did all of the interviews  
15  
16 and data collection.  
17  
18

19  
20 **Data sharing statement:** All available data can be obtained by contacting the corresponding  
21  
22 author.  
23  
24

25  
26 **Word count: 4,479**  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## Abstract

**Rationale:** Physical rehabilitation (encompassing early mobilisation) of the critically ill patient is recognized best practice, however further work is needed to explore the patients' experience of rehabilitation qualitatively; a better understanding may facilitate implementation of early rehabilitation, and elucidate the journey of survivorship.

**Objectives:** To explore patient experience of physical rehabilitation from critical illness during and after a stay on ICU.

**Design:** Exploratory grounded theory study using semi-structured interviews.

**Setting:** Adult medical/surgical ICU of a London teaching hospital.

**Participants:** A purposive sample of ICU survivors with intensive care unit acquired weakness (ICUAW) and an ICU length of stay of >72 hours.

**Analysis:** Data analysis followed a four-stage constant comparison technique: open coding, axial coding, selective coding, and model development, with the aim of reaching thematic saturation. Peer debriefing and triangulation through a patient support group were carried out to ensure credibility.

**Main results:** Fifteen people were interviewed (with four relatives in attendance). The early rehabilitation period was characterized by episodic memory loss, hallucinations, weakness, and fatigue, making early rehabilitation arduous and difficult to recall.

Participants craved a paternalised approach to care in the early days of ICU.

The central idea that emerged from this study was recalibration of the self. This is driven by a lost sense of self, with loss of autonomy and competence; dehumanized elements

1  
2  
3 of care may contribute to this. Participants described a fractured life narrative due to  
4  
5 episodic memory loss, meaning that patients were shocked on awakening from sedation  
6  
7 by the discrepancy between their physical form and cognitive representation of  
8  
9 themselves.  
10  
11  
12

13  
14 **Conclusions:** Recovery from ICUAW is a complex process that often begins with  
15  
16 survivors exploring and adapting to a new body, followed by a period of recovering  
17  
18 autonomy. Rehabilitation plays a key role in this recalibration period, helping survivors  
19  
20 to reconstruct a desirable future.  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52

53 **Key words:** critical care, early mobilisation, rehabilitation, patient experience, recovery,  
54 physical therapy.  
55

56 **Abstract word count:** 298  
57  
58  
59  
60

## Strengths and Limitations of this study

1. This was an exploratory qualitative grounded theory study using semi structured interviews with survivors of critical illness to explore their experience of physical rehabilitation after critical illness; the approach adopted, and the data generated provided an extremely rich source of individual experience with many consistent features.
2. A constant comparison technique of data analysis was used, and enrolment continued until thematic saturation was reached.
3. Triangulation and peer debriefing were completed to ensure credibility of the study findings that clearly resonated with an independent group of critical illness survivors.
4. The patients were all recruited from one centre, which may limit transferability of findings. Qualitative studies of this kind innately have a small sample size, however, the richness of the data produced allows deep exploration of meaning and model development and thematic saturation was also reached.
5. The variation in time to interview may be considered a limitation of this study in view of impaired recall for longer gaps, however, there was no notable difference in the richness of memories and insight provided by those interviewed at different time points. The variation in time to interview also elucidated the process of recovery over time.

## Introduction:

Rapid muscle wasting<sup>1,2</sup>, functional decline<sup>3</sup> and delirium<sup>4</sup> are common consequences of critical illness. In the long term, they can lead to prolonged periods of weaning from mechanical ventilation, disability, reduced endurance, anxiety, and depression<sup>5-9</sup>. To combat these issues mobilisation, minimising sedation and spontaneous breathing should be instigated early, with research demonstrating safety and likely efficacy<sup>10-13</sup>. Furthermore, on-going rehabilitation following discharge from critical care, and attendance at ICU follow up clinics are also advocated, although research showing direct benefit of these interventions is limited<sup>14-15</sup>.

Although implementation of early mobilisation protocols and post-ICU rehabilitation is inconsistent<sup>14,16-17</sup>, early adopters of these strategies are striving for them to become the norm<sup>12,13,18</sup>. In such centres, it is not uncommon for patients to receive active out of bed physical rehabilitation whilst receiving full mechanical ventilation, renal replacement therapy, and inotropic support<sup>18</sup>. Due to the severity of weakness that can be associated with prolonged critical illness, these rehabilitation sessions are often delivered by two or more therapists/nurses, and can require technical equipment and physical handling. It is perhaps unsurprising that pain, fatigue, weakness, anxiety, fear, lack of motivation and patient confidence are reported as barriers and reasons for cessation of early rehabilitation<sup>19, 20</sup>.

Sottile and colleagues (2015)<sup>21</sup> completed a survey of patient experience of early mobilisation in ICU concluding that patients recognized the importance of early mobilisation, but found it difficult, tiring and uncomfortable.

1  
2  
3 In spite of current enthusiasm, there is a paucity of literature exploring survivors'  
4 experience of early mobilisation and physical rehabilitation during and after a stay in ICU  
5  
6 in an in-depth manner. For the purpose of this manuscript, the term 'rehabilitation' is  
7  
8 used to encompass early mobilisation and physical rehabilitation implemented by  
9  
10 physiotherapists from admission to ICU.  
11  
12  
13  
14  
15

16 **Aim:** To explore the patient experience of recovery from critical illness, with emphasis  
17  
18 on their experience of rehabilitation, and to develop a theoretical model grounded in  
19  
20 these data.  
21  
22

23  
24 **Methods:**

25  
26  
27 **Qualitative approach and research paradigm:** Constructivist grounded theory study<sup>22</sup>  
28  
29 using semi-structured interviews with a purposive sample of adult ICU survivors.  
30  
31 Constructivism contends that individuals' views are directly influenced by their  
32  
33 experiences, and it is these individual experiences and views that shape their perspective  
34  
35 of reality. Constructivists believe that individuals have different realities that will be  
36  
37 influenced by context- this is a 'relativist' ontological stance<sup>22</sup>.  
38  
39  
40  
41  
42

43 Constructivist grounded theory is an appropriate methodology for this study because it  
44  
45 allows the researcher to develop a theoretical model to explain the data based on an  
46  
47 iterative process of data immersion, analysis and interpretation, which recognises and  
48  
49 accounts for contextual factors.<sup>22</sup>  
50  
51  
52

53 **Setting:** Participants were recruited from the adult medical/surgical ICU of a 430 bedded  
54  
55 London teaching hospital between November 2015 and September 2016.  
56  
57  
58  
59  
60



1  
2  
3 **Participants and sampling:** Participants were purposively sampled. Screening and  
4 inclusion criteria were: English-speaking, a critical stay of >72 hours, capable of providing  
5 informed consent determined using the Mental Capacity Act assessment<sup>23</sup>, anticipated  
6 to survive, aged over 18 and documented ICUAW determined via case note review (this  
7 was to ensure that the participants had exposure to rehabilitation interventions.)  
8  
9

10  
11  
12  
13  
14  
15  
16 The clinical team identified potential participants against the broad inclusion criteria  
17 stated above to ensure that it was appropriate for them to be approached by the  
18 research team. Notes were screened with the aim of purposively selecting a varied  
19 sample of participants that could speak to the breadth of emerging themes. If deemed  
20 appropriate participants were then approached by the lead researcher (EJC) and  
21 provided with written information. If they had capacity to consent and agreed to  
22 participate, written informed consent was gained. Participants that could not provide  
23 informed consent were excluded.  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35

36 At the discretion of the participant, relatives were also invited to be present in the  
37 interview to enable exploration and elucidation of any ICU-associated memory loss. As  
38 the study progressed participants were selected to ensure a heterogeneous sample, with  
39 the aim of achieving thematic saturation<sup>22,24-26</sup>, for example, targeting varying degrees of  
40 ICUAW, different genders, and specific age categories.  
41  
42  
43  
44  
45  
46  
47  
48

49 **Ethics:** This study was granted approval by the East of England Ethics committee (REC  
50 reference number 14/EE/1027) and from the Research and Development Department at  
51 the study site.  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 **Data collection methods:** The semi-structured interviews were conducted by EJC. For  
4  
5 reflexivity, EJC is research physiotherapist with expertise in critical care and prior training  
6  
7 in qualitative methods research. EJC also has personal experience of major injury and as  
8  
9 a close relative of an ex ICU patient. SJB is an ICU medical consultant and is involved with  
10  
11 ICU follow up clinics, and EJM is a researcher focusing on management and change in the  
12  
13 health sector, with expertise in qualitative methodology.  
14  
15  
16

17  
18 The interviews followed a topic guide designed with input from the Intensive Care  
19  
20 Society Patient and Relatives Group (Table 1). The questions in the topic guide focused  
21  
22 on the memory of the admission to ICU, any physical weakness that they encountered,  
23  
24 and patients' experience of rehabilitation in the ICU and following discharge. The  
25  
26 questions were intentionally left open to initiate reflections and to allow subsequent  
27  
28 detailed exploration of the issues that appeared important to the interviewee. The first  
29  
30 interview was used as a pilot, however, as no changes were made and those data  
31  
32 collected from this interview were rich and informative, it was retained and analysed in  
33  
34 the results.  
35  
36  
37  
38  
39

40  
41 As the study progressed and themes emerged, participants were asked to elaborate and  
42  
43 probed on specific issues in line with the constant comparison technique, for example:  
44  
45 how the perception of the physiotherapist's strength influenced their rehabilitation  
46  
47 experience; how they perceived their body now; what differences there were between  
48  
49 their current and previous physical function; how they saw their future; and what they  
50  
51 defined as physical rehabilitation.  
52  
53  
54

55  
56 The interviews were carried out either in the hospital or in the community after ICU  
57  
58 discharge. Enrolment and interviews continued until thematic saturation was reached  
59  
60

1  
2  
3 i.e. no new ideas were emerging, as per the criteria outlined by Bonde (2013)<sup>26</sup>. This was  
4  
5 to challenge the emergent model and ensure credibility. The interviews were  
6  
7 anonymised, recorded, and transcribed *verbatim* by a professional transcription  
8  
9 company. All transcripts were double-checked for accuracy by EJC.  
10  
11  
12

13  
14 All interviewees were given pseudonyms to ensure anonymity. Further demographic and  
15  
16 clinical data were also collected from the case notes: age, critical care and hospital  
17  
18 length of stay, APACHE II score, admission diagnosis, residence prior to admission, pre-  
19  
20 morbid functional level, and hospital discharge destination.  
21  
22

23  
24 **Data processing and analysis:** Transcripts were uploaded onto Nvivo® software (QSR  
25  
26 International, Doncaster, Australia) for analysis. They were read and reread by EJC to  
27  
28 ensure full immersion in the data. Memo writing was used throughout. The first stage of  
29  
30 the analysis process is 'open coding', which is the identification of primary broad  
31  
32 categories; these may be around a theme or topic, or more conceptual, such as emotion  
33  
34 or attitude. The second stage is 'axial coding'; here categories are clustered together into  
35  
36 meaningful, related groups. The third stage is 'selective coding', where core themes are  
37  
38 identified. Lastly, the themes are used to generate a theoretical framework to explain  
39  
40 the data<sup>22,24-26</sup>. Data collection and analysis occur concurrently, so that constant  
41  
42 comparison was made between emerging themes (both within and between narratives),  
43  
44 and the literature, allowing model refinement. After the fourth interview had been  
45  
46 transcribed and open coding had been completed, axial codes began to form. These  
47  
48 ideas were then discussed in detail with the research team. This was followed by a  
49  
50 dynamic process of reflection after each interview to develop and refine the axial codes  
51  
52 into selective codes until a model encompassing all elements was developed. The last  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 interviews were used to challenge this model and to assess for data saturation. This  
4  
5 process allowed a central phenomenon to emerge from the data<sup>22,24-26</sup>. The words used  
6  
7 for coding were based on the lead researchers interpretation and terms in related  
8  
9 literature.  
10  
11  
12

### 13 14 **Techniques to enhance trustworthiness:**

#### 15 16 17 **Peer debriefing:**

18  
19  
20 Peer debriefing was completed via in-depth discussion with SJB and EJM. This was done  
21  
22 regularly throughout the course of the study.  
23  
24

#### 25 26 **Patient and public involvement**

27  
28  
29 A patient representative from an ICU support group was consulted in the development  
30  
31 of the topic guide. An initial draft of the topic guide was developed by the research team  
32  
33 and it was then sent to the patient representative for review and modification, all of  
34  
35 their recommended changes were made. Patients and public were not involved in the  
36  
37 recruitment or conduct of the study. Participants were given the opportunity to receive  
38  
39 information on the results of the study at their request.  
40  
41  
42  
43

#### 44 45 **Triangulation**

46  
47  
48 Triangulation and sense checking was completed through presentation to an ICU  
49  
50 survivor support group with subsequent dialogue to assess the dependability,  
51  
52 confirmability and credibility of the model (this did not include interview participants).  
53

54  
55  
56 At the support group the model was presented and then there was opportunity for  
57  
58 questions and answers with the researcher (EJC). The group were then left to discuss the  
59  
60

1  
2  
3 study between themselves and feedback to the research team with any thoughts at a  
4  
5 later date to allow them to speak openly and frankly with each other. The leader of the  
6  
7 support group fed back to the research team that the participants resoundingly agreed  
8  
9 with the concepts and felt that the work '*encompassed all the areas that were important*  
10  
11 *and relevant to those who have experienced critical illness*'. The presentation also led to  
12  
13 a very tearful response from some attendees who reported to have felt 'understood'.  
14  
15  
16  
17

### 18 **Results:**

19  
20  
21 Eleven hours of qualitative data from 15 participants (with 4 additional relatives present)  
22  
23 were collected. The *patients* are described in Table 2.  
24  
25

26  
27  
28 *[Insert table 2: Patient demographics]*  
29

30  
31 The median ICU and hospital length of stay were 19 days (IQR 8-33) and 63 days (IQR 34-  
32  
33 107) respectively. The median time between ICU discharge and interview was 56 days  
34  
35 (IQR: 36-80). Ten (66.6%) of the interviews took place at the hospital whilst the patients  
36  
37 were still inpatients, and five (33.3%) took place after discharge in the patient's home  
38  
39 (n=2), work (n=1), or in a clinic room (n=2). The interviews lasted a median of 39  
40  
41 minutes (IQR: 28-50).  
42  
43  
44  
45

46  
47 The central phenomenon grounded in these data was *recalibration of the self*. There  
48  
49 were two themes contributing to this temporal model of recovery: the transition '*from*  
50  
51 *prior self to current self*', and the transition '*from current self to construction of the*  
52  
53 *future self*'. When questioned about early physical function, patients recalled a  
54  
55 discrepancy at the time of recovering awareness between their *current self*, which  
56  
57 incorporates their physical dependency, fatigue, clarity of mind, and self-image, and the  
58  
59  
60

1  
2  
3 mental representation of themselves, which is still consistent with their *preadmission*  
4  
5 *self*. This discrepancy seemed to be due to episodic memory loss of their admission  
6  
7 period. Additionally, patients lacked some of the cognitive requirements for prospection  
8  
9 at this point<sup>26</sup> and therefore they struggled to envisage a compelling future self. This  
10  
11 appears to lead on to a period of *recalibration*.  
12  
13  
14  
15

16 Although this central phenomenon of *recalibration* may seem distinct from the early  
17  
18 physical rehabilitation experience that was the focus of this study, it was quite the  
19  
20 opposite, with the process of *recalibration* seeming inextricably linked to the  
21  
22 rehabilitation experience. Physical independence and function are core components of  
23  
24 the concept of *self*. When physical ability deteriorates so unexpectedly, rapidly, and  
25  
26 without obvious causation (as in ICUAW) it comes as a shock to the patient blurring their  
27  
28 sense of self. Physical rehabilitation aims to improve impairments and function by  
29  
30 challenging patients' physical ability thereby, in this extreme context, inadvertently  
31  
32 challenging their self-perception as well.  
33  
34  
35  
36  
37  
38

39 This model suggests that physical rehabilitation within ICU helps patients to challenge  
40  
41 and explore their current functional level and reconcile their self-discrepancy i.e.  
42  
43 difference between their physical self and the cognitive image of themselves. The  
44  
45 process of therapy goal-setting also challenges their capacity to think about the future;  
46  
47 discussing goal setting with participants therefore elucidated the difficulties they may  
48  
49 have in constructing a compelling future to act as a motivational force.  
50  
51  
52  
53

54 The rationale and contributing themes are presented below. Supporting evidence is  
55  
56 presented in Table 3.  
57  
58  
59  
60

## ***'From prior self to current self'***

### **Episodic memory loss**

Fundamental to the patient experience of rehabilitation, and underpinning the theoretical interpretation, was patients' episodic memory loss (i.e. loss of a specific autobiographical event) of their admission to ICU, regardless of their admission background or diagnosis. In some cases this memory gap lasted weeks, with some participants unable to recall any rehabilitation sessions on ICU at all, citing their ward rehabilitation sessions as their first experiences. The first clear memory *for all* participants was a family member at the bedside. This frequently elicited a tearful response, for example, George stated: "*(my first memory on awakening) was my mother stroking my arm, saying 'Mum's here'... that was some 30 days after my admission*". This memory loss is of paramount importance, as it made it difficult for participants to rationalise and understand their current situation.

### **Hallucinations and delusions**

All patients experienced vivid hallucinations that often involved torture and trying to escape some of the hallucinations, however, were pleasant experiences, such as a friendly dog in the ICU. Those with a history of recreational drug use seemed less shocked by hallucinations and able to rationalise their mental state, for example, John stated: "*you're pumped full of so many drugs, it doesn't surprise me that you're tripping out.*" Perceived stigma influenced patients' comfort in discussing hallucinations with

1  
2  
3 staff, for example, EJC asked Tom: *“Did you tell anyone about the hallucinations at the*  
4  
5 *time?”* Tom replied: *“No... I just felt a bit silly”*.  
6  
7  
8  
9  
10

## 11 **Weakness**

12  
13  
14  
15 On awakening patients reported frustration at their inability to communicate and were  
16  
17 shocked by the severity of their weakness, as mentally they still saw themselves as  
18  
19 capable of the physical tasks they were able to do pre-admission, for example, Sasha  
20  
21 stated: *“I didn't realise I couldn't walk. I thought I could and I tried to get out of bed loads*  
22  
23 *of times, but up here I was weak (legs) and the top of my arms were weak as well. I*  
24  
25 *couldn't do it.”* Their actual *physical-self* and cognitive representation of themselves did  
26  
27 not match, for example, Sarah said: *“I just happened to catch sight of my whole body (in*  
28  
29 *the mirror) and I nearly died. I thought; ‘that doesn't resemble the person that I am’.”*  
30  
31 However, it was the psychological symptoms that were of the greatest concern to  
32  
33 patients initially, for example, Evan felt that: *“there were tubes all over the place... but*  
34  
35 *that was the least of my worries. The specialists were there, and my son. I said ‘I don't*  
36  
37 *know who that is (son)’. My son came back on the Monday, then I recognised him and*  
38  
39 *things started falling back into place.”*  
40  
41  
42  
43  
44  
45  
46  
47

## 48 **“Noxious cycle” of ICU**

49  
50  
51 Overwhelming fatigue, insomnia (due to noise and disruption), boredom and the  
52  
53 inability to concentrate were prevalent, which had a negative impact on the ability to  
54  
55 engage in both physical rehabilitation and cognitive tasks, and made many fear early  
56  
57 rehabilitation, for example, John stated: *“Physically tiring, emotionally, you're like “sh\*t,*  
58  
59  
60



1  
2  
3 *really? I've got to do it (physiotherapy) now. I haven't got any energy at all."* Procedural  
4  
5 pain was reported in only a few instances, but discomfort was problematic, for example,  
6  
7 from being 'swaddled' in blankets (John). For many this seemed to form a "noxious  
8  
9 cycle" (Figure 1).  
10  
11  
12  
13  
14  
15  
16

### 17 **Humanisation of care**

18  
19  
20 Participants valued "humanised" care<sup>27</sup>, often remembering the staff members who  
21  
22 made them laugh and feel safe, for example, Caroline said: "*I remember one bloke, one*  
23  
24 *nurse, who- he would come in and smile, and I said 'Oh, you're always smiling. You make*  
25  
26 *me so happy'."* Trust in the clinical team was also important; if trust was compromised  
27  
28 then it had a negative impact on participants engagement with rehabilitation. Trust  
29  
30 seemed dependent on the rapport the staff member developed with the patient,  
31  
32 including their ability to communicate honestly and to maintain patient's hope, for  
33  
34 example, Michelle stated: "*I trust him... because when Tom (physiotherapist) says*  
35  
36 *something, it's true. Everything he said was true."* However, the staff-patient interaction  
37  
38 was not always positive, with many patients describing examples of *de-humanised*  
39  
40 care<sup>27</sup>. This included loss of agency: "*I feel so not free, everyone is doing what they want,*  
41  
42 *I'm like a puppet and I hate that"* (Michelle); and feeling isolated: "*I don't think I had a*  
43  
44 *voice at one point, which was probably one of the most difficult things to experience,*  
45  
46 *because you can't talk to people"* (Richard).  
47  
48  
49  
50  
51  
52  
53

54  
55 Although not related to humanisation of care, the physical attributes of staff also  
56  
57 influenced patients' rehabilitation experience, if physiotherapists looked small, young,  
58  
59  
60

1  
2  
3 and weak, then patients had less trust in their physical ability to keep them safe during  
4 rehabilitation, an example came from David: *“He (the physio) was strong of course. One*  
5  
6  
7  
8 *admires that. It’s an ability, you know; and of course, not everybody’s going to have that*  
9  
10 *ability.”*

### 14 ***‘From current self to construction of the future self’***

#### 17 **Recognising milestones to recovery and goal setting**

20 The vulnerability described by patients and relatives seemed to lead to a sense of  
21  
22  
23 desired paternalism in the early days; they did not feel ready to be in control. This was  
24  
25 further perpetuated by a lack of understanding of the stages of recovery, for example,  
26  
27 Tom stated: *“how you are improving may not be quite so obvious to the patient”*. The  
28  
29 memory loss of their admission meant that patients did not recall their acute stages of  
30  
31  
32 illness, and hence their physical decline; the weakness that they were experiencing did  
33  
34 not make sense, and was often so severe that it made it difficult to envisage the next  
35  
36  
37 steps in their recovery. As a result, patients did not always recognise basic functional  
38  
39  
40 tasks as rehabilitation or indeed their achievements as progress, for example, Michelle  
41  
42 stated: *“The other day the whole ward congratulated me- and even now I feel*  
43  
44 *embarrassed – because I washed myself. I didn’t wait until now to know how to wash*  
45  
46 *myself; I thought it was so stupid.”* Therapeutic adjuncts, such as the use of a bed bike  
47  
48 or tilt table, were more commonly recalled as rehabilitation.

51  
52  
53 As patients had limited understanding of the recovery milestones early on, they wanted  
54  
55 the multi-disciplinary team to set their rehabilitation goals as *“they did not know what*  
56  
57 *goals to set”* (Jim). The main thing that kept them focused on engaging in rehabilitation  
58  
59  
60

1  
2  
3 at this point was their family and loved ones, Sarah described this: "I cry a  
4  
5 lot...something helped me to keep going, an inner strength came...the kids..."  
6  
7

8  
9 Patient involvement in early goal setting was described as like "*being in a car crash and*  
10  
11 *someone asking you how you want to be cut out.*" Most patients had a 'just get on with  
12  
13 it' approach to rehabilitation. Martin: "*I just blind folded said, 'if this is what I am*  
14  
15 *supposed to do, I will do it.*" However, despite desiring early clinician-led rehabilitation,  
16  
17 all patients identified a high-level goal that aligned to the core values of who they are;  
18  
19 examples include, returning to work, going on holiday, finishing a PhD, and getting  
20  
21 married.  
22  
23  
24

25  
26 As patients progressed through the stages of recovery, they started to recognise smaller  
27  
28 functional gains as improvement and engaged more in the goal-setting and  
29  
30 rehabilitation planning process, for example, Sarah stated: "*Well, I was shocked at how*  
31  
32 *little I could do, but now, it's the other way, I'm actually shocked at how much I can do*  
33  
34 *and I am doing. It's really good.*" Their yardstick for comparison now became who they  
35  
36 were on awakening, and not who they were prior to admission; they were recalibrating.  
37  
38  
39  
40

## 41 42 **Discussion**

43  
44  
45 This work focused on exploring the experience of physical rehabilitation after critical  
46  
47 illness, however, as with inductive research, what transpired was a complex model of  
48  
49 recovery extending beyond the physical. Patients demonstrated an interruption to  
50  
51 personal narrative, a lost sense of self associated with loss of autonomy, temporary  
52  
53 desired paternalism and gave examples of accidental dehumanised care (albeit mostly  
54  
55 non-maleficent in intent). Delirium, sleep deprivation, fatigue and memory loss acted as  
56  
57  
58  
59  
60

1  
2  
3 potent mediators between the patients' physical impairments, and their ability to  
4  
5 recalibrate to their new disability, and engage in rehabilitation.  
6  
7

8  
9 It is interesting to consider these findings in the context of established psychological  
10  
11 theory. Deci & Ryan's Self-Determination Theory<sup>28,29</sup> attempts to explain why people  
12  
13 engage in goal-orientated behaviour, exploring how this leads to well-being and personal  
14  
15 growth. Its three core concepts are: *autonomy* (the ability to be in control of oneself),  
16  
17 *competence* (the ability to manage the situation they are in) and *relatedness* (the ability  
18  
19 to have an emotional connection with others). Only when these needs are met can  
20  
21 intrinsic motivation flourish. Critical illness can strip patients of *autonomy* and  
22  
23 *competence*, and perhaps for a shorter period, *relatedness*. In the initial stages of critical  
24  
25 illness, patients may be unable to communicate and talk, be unable to move easily due  
26  
27 to weakness, may have hallucinations, and be too fatigued to engage in decision making.  
28  
29 Hence, loss of *autonomy* and *competence* are key features of the patient experience.  
30  
31 *Relatedness*, which may recover earlier (or be encouraged) is of paramount importance  
32  
33 to them, and was a motivator to engage in rehabilitation.  
34  
35

36  
37 Markus and Nurius (1986) developed a theory called "The Possible Self"<sup>30</sup>. They contend  
38  
39 that humans have different cognitive representations of who we are (*current self*) and  
40  
41 who we could be (*possible self*). *Possible selves* drive behaviour. A notion of the *possible*  
42  
43 *self* helps us to assess our *current self* by creating a benchmark for comparison for self-  
44  
45 evaluation. It can also provide tangible rehabilitation goals. *The possible self* and goal  
46  
47 setting also rely heavily on temporality, therefore requiring narrative of the past, and the  
48  
49 capacity to prospect. Physiologically, prospection depends on episodic memory,  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 prospective memory, emotional stability and hypothetical thinking<sup>31</sup>, some of which can  
4  
5 be impaired in critical illness due to sleep deprivation, fatigue and delirium.  
6  
7

8  
9 On awakening, patients' immediate cognitive representation of their *current self*  
10  
11 matched their pre-admission self because they do not remember their functional  
12  
13 decline, however, their body had changed. Their mental image of their *current self* and  
14  
15 their *physical self* were not aligned. Furthermore, patients could not remember the  
16  
17 totality of their past, they did not recognise their present, and they struggled to  
18  
19 construct a compelling future self. This impaired their ability to engage in rehabilitation  
20  
21 goal setting and led to a sense of vulnerability, desired paternalism and emphasis on  
22  
23 *relatedness*.  
24  
25  
26  
27

28  
29 This model of *recalibration* ties these established psychological ideas together, reflecting  
30  
31 the need for patients to explore their new self, adapt to it and allow it to become their  
32  
33 new yardstick. When this was achieved, smaller milestones in recovery became  
34  
35 meaningful goals. Others have described similar concepts as a *liminal* state. Liminality is  
36  
37 an anthropological term from the Latin word *līmen*, meaning threshold<sup>32</sup>. It refers to  
38  
39 someone who is transitioning. It is often associated with a change in role/identity, or a  
40  
41 loss of one self to be replaced with another. This can create inner turmoil, especially if  
42  
43 that change is not invited.  
44  
45  
46  
47

48  
49 This idea of liminality in ICU has been touched upon by a number of authors<sup>32-35</sup>. Kean  
50  
51 and colleagues<sup>32</sup> identified 'unscheduled status passage' from *prior self* to *critically ill self*  
52  
53 as a theme in a longitudinal study of ICU survivorship. They found that this unscheduled  
54  
55 liminal stage is worsened by memory loss and delirium, and that this process of change  
56  
57  
58  
59  
60

1  
2  
3 is temporal in nature, both progressing and regressing (in the event of decline). In order  
4  
5 to move on, patients need to regain autonomy.  
6  
7

8  
9 Lindberg and colleagues<sup>35</sup> described what recovering autonomy looks like, suggesting  
10  
11 that patients go through four stages: the first is to acknowledge their dependence (or  
12  
13 paternalism), and then they strive to be recognised as a person (“humanised” care).  
14  
15 These two stages echo the findings of this study. The latter two stages are ‘invited  
16  
17 participation in care’, and ‘becoming a co-partner in the decision-making’. These stages  
18  
19 describe how staff coach patients to take control again through mutual trust,  
20  
21 understanding and co-determination.  
22  
23  
24

25  
26 Although these may seem like abstract concepts, it is the authors’ view that they have  
27  
28 direct relevance to clinical practice, especially as early rehabilitation becomes a key  
29  
30 aspect of acute care. The reason for this is that perception of self and engagement in  
31  
32 rehabilitation and goal-setting are inextricably linked.  
33  
34  
35

36  
37 If a person’s mind is telling them one thing about who they are and what they are  
38  
39 capable of doing, and their body is telling them another, they cannot start thinking about  
40  
41 the future until they reconcile that difference. They cannot reconcile that difference with  
42  
43 delirium, hallucinations and lack of episodic memory to justify their current situation and  
44  
45 facilitate hypothetical thinking and propection<sup>31</sup>. Combining psychological intervention  
46  
47 with physiotherapy intervention may help to address this.  
48  
49  
50

51  
52 Goal-setting depends on the capacity to prospect. It is also a key recommendation in the  
53  
54 UK National Institute for Health and Care Excellence (NICE) Guidelines for Rehabilitation  
55  
56 after Critical Illness<sup>36</sup> and NICE Quality Standards<sup>37</sup>. The Quality Standards state that  
57  
58  
59  
60

1  
2  
3 rehabilitation goals should be set within 4 days of admission, and ideally should be  
4 patient-agreed. These data would suggest that asking *patients* to set goals at day 4 may  
5  
6 be premature. Further research exploring the application of the model of recovering  
7  
8 autonomy described by Lindberg<sup>35</sup> may assist in tailoring rehabilitation guidelines to the  
9  
10 specific needs of the critically ill.  
11  
12  
13  
14

15  
16 Further focus on *how* rehabilitation is delivered, not just *what* is delivered could also be  
17  
18 instructive. The impact of the dynamic between a sports coach and the players is well  
19  
20 known, yet this coaching dynamic is neglected somewhat in ICU rehabilitation. If  
21  
22 clinicians are able to assist patients in recalibrating to their *new current self*, and the  
23  
24 reconstruction of a compelling *future self*, it may improve patient care and outcome.  
25  
26 Further research will be needed to confirm the concepts identified in this initial  
27  
28 exploratory study. However, we believe the concepts identified are sufficiently plausible  
29  
30 and robust to pose challenges to clinicians working with recovering critically ill patients  
31  
32 (outlined in Box 1).  
33  
34  
35  
36  
37  
38

39 **Box 1: Key observations and challenges to practice.**

41 **Key observations**

- 42  
43 ❖ Patients recovering consciousness during or after a critical illness are likely to be  
44 shocked by the transition through which they have gone. Part of that shock is the  
45 due to the unplanned interruption of their autobiographical story.  
46  
47 ➤ How can you help to fill the gaps in autobiographical memory?  
48  
49  
50  
51  
52 ❖ Patients' immediate memory is of who they were and what they were able to do  
53 before their critical illness. This is in collision with what they can actually do and a  
54 period of recalibration is needed to allow people to align the two and develop  
55 reasonable ambitions and goals.  
56  
57 ➤ How can you support patients to explore their current function and settle the  
58  
59  
60

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

discrepancy between expectations and reality?

- ❖ This recalibration is the development of an understanding of the relationships between their past, present and possible futures selves.
  - How can you help patients to envisage a compelling future?
- ❖ Because of this need for recalibration along with delirium and impaired cognition patients may need, and wish for, assistance in planning early rehabilitation. As autonomy recovers, patients desire to become fully involved increases.
  - How can you recognise and support recovering autonomy?
- ❖ Motivation and engagement are crucial in maximising the benefits of rehabilitation. Leveraging human relationships (relatedness) and encouraging autonomy are likely to be helpful; care that is de-humanising, even if “efficient” is likely to impair recovery.
  - How can humanisation of care be optimised in your ICU?

**Acknowledgements:** Thanks to Dr Cherry Kilbride and Dr William Young (Brunel University London) for their input into the theoretical interpretation of these data. Thank you to Gordon Sturmey and the ICU support network in Reading for their involvement and support with the study.



**References:**

1. Puthuchearry ZA, Rawal, JR, McPhail M, Connolly B, Ratnayake G, Chan P, Hopkinson NS, Phadke R, Dew T, Sidhu PS, Velloso C, Seymour A, Agley CC, Selby A, Limb M, Edwards LM, Smith K, Rowlerson A, Rennie MJ, Moxham J, Harridge S, Hart N, Montgomery HE. Acute Skeletal Muscle Wasting in Critical Illness. *JAMA*, 2013; 310 (15), 1591-1600.
2. Friedrich O, Reid MB, Van den Berge G, Vanhorebeek I, Hermans G, Rich MM, Larsson L. The sick and the weak: Neuropathes/myopathies in the critically ill. *Physiol Rev*. 2015; 95 (3), 1025-109.
3. Hermans G, & Van den Berghe G. Clinical review: intensive care unit acquired weakness. *Crit Care*. 2015; 5 (19), 274.
4. Cavalazzi R, Saad M, Marik PE. Delirium in the ICU: an overview. *Ann Intensive Care*. 2012; 2 (49).
5. Kress JP & Hall JB. Intensive care unit acquired weakness and recovery from critical illness. *N Eng J Med*. 2014; 370, 1626-1635.
6. Cuthbertson BH, Elders A, Hall S, Taylor J, MacLennan G, MacKirdy F, Mackenzie SJ, and the Scottish Critical Care Trials Group and the Scottish Intensive Care Society Audit Group. Mortality and quality of life in the five years after severe sepsis. *Critical Care*. 2013; 17, R70.
7. Kaukonen K-M, Bailey M, Suzuki S, Pilcher D, Bellomo R. Mortality related to severe sepsis and septic shock among critically ill patients in Australia and New Zealand, 2000-2012. *JAMA*. 2014; 311 (13), 1308-1316.
8. Hill AD, Fowler RA, Pinot R, Herridge MS, Cuthbertson BH, Scales DC. Long-term outcomes and healthcare utilization following critical illness – a population-based study. *Critical Care*. 2016; 20, 76.
9. Pandharipande P, Girard TD, Jackson JC, Morandi A, Thompson JL, Pun NE, Brummel CG, Highes EE, Vasilevskis AK, Shintani KG, Moons SK, Geevaghese A, Canonico RO, Hopkins RO, Bernad GR, Dittus RS, Ely EW. Long-Term Cognitive Impairment after Critical Illness. *N Engl J Med*. 2013: 369, 1306-1316.
10. Nydahl P, Sricharoenchai T, Chandra S, Kundt FS, Huang M, Fischill M., Needham D.M. Safety of Patient Mobilization and Rehabilitation in the Intensive Care Unit.

- 1  
2  
3 Systematic Review with Meta-Analysis. *Annals of the American Thoracic Society*.  
4 2017; 14 (5).  
5  
6  
7 11. Castro-Avila AC, Seron P, Fan E, Gaete M, Mickan S. (2015) Effect of early  
8 rehabilitation during Intensive Care Unit stay on functional status: systematic review  
9 and meta-analysis. *PLoS ONE*. 2015; 10 (7), e0130722.  
10  
11 12. Balas MC, Burke WJ, Gannon D, Cohen MZ, Colburn L, Bevil C, Franz D, Olsen KM, Ely  
12 WE, Vasilevski EE (2013) Implementing the awakening and breathing coordination,  
13 delirium monitoring/management, and early exercise/mobility bundle into everyday  
14 care: opportunities, challenges, and lessons learned from implementing the ICU  
15 pain, agitation and delirium guidelines. *Crit Care Med*. 2013; 41: S116-S127.  
16  
17 13. Balas MC, Vasilevskis EE, Olsen KM, Schmid KK, Shostrum V, Cohen MZ, Peitz G,  
18 Gannon DE, Sisson J, Sullivan J, Stohtert JC, Lazure J, Nuss SL, Jawa RS, Freihurt F, Ely  
19 EW, Burke WJ. (2014) Effectiveness and safety of the awakening and breathing  
20 coordination, delirium monitoring/management, and early exercise/mobility bundle.  
21 *Crit Care Med*. 2014; 42:1024-1036.  
22  
23 14. Connolly B, Douiri A, Steier J, Moxham J, Denehy L, Hart N. A UK survey of  
24 rehabilitation following critical illness: implementation of NICE Clinical Guidance 83  
25 (CG83) following hospital discharge. *BMJ Open*. 2014; 4; e004963.  
26  
27 15. Jensen JF, Thomsen T, Overgaard D, Bestle MH, Christensen D, Egerod I. Impact of  
28 follow-up consultations for ICU survivors on post-ICU syndrome: a systematic review  
29 and meta-analysis. *Intensive Care Med* (2015) 41:763–775  
30  
31 16. Nydahl P, Parker RA, Bartoszek G, Dubb R, Filipovic,S, Flohr H-J, Kaltwasser A, Mende  
32 H, Rothaug O, Schuchhardt D, Schwabbauer N, Needham D. Early Mobilization of  
33 Mechanically Ventilated Patients: A 1-Day Point-Prevalence Study in Germany.  
34 *Critical Care Medicine*. 2014; 42 (5), 1178-1186.  
35  
36 17. Berney SC, Rose JW, Bernhardt J, Denehy L, Prospective observation of physical  
37 activity in critically ill patients who were intubated for more than 48 hours. *Journal*  
38 *of Critical Care*. 2015; 30(4), 658-663.  
39  
40 18. Hodgson CL, Stiller K. Needham D, Tipping CJ, Harrold M, Baldwin CE, Bradley S,  
41 Berney S, Caruana LR, Elliott D, Green M, Haines K, Higgins AM, Kaukonen KM,  
42 Leditschke IA, Nickels MR, Paratz J, Patman S, Skinner EH, Young PJ, Zanni JM,  
43 Denehy L, Webb SA. Expert consensus and recommendations on safety criteria for  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

- 1  
2  
3 active mobilisation of mechanically ventilated critically ill patients. *Critical Care*.  
4 2014; 16:658.  
5  
6  
7 19. Parry S, Knight LD, Connolly B, Baldwin C, Puthuchery Z, Morris P, Mortimore J, Hart  
8 N, Denehy L, Granger CL. Factors influencing physical activity and rehabilitation in  
9 survivors of critical illness: a systematic review of quantitative and qualitative  
10 studies. *Intensive Care Med*. 2017; 43, 531–542.  
11  
12  
13  
14 20. Wright SE, Thomas K, Watson G, Baker C, Bryant A, Chadwick, TJ, Shen J, Wood R,  
15 Wilkinson J, Mansfield L, Stafford V, Wade C, Furneal J, Henderson A, Hugill K,  
16 Howard P, Roy A, Bonner S, Baudouin S. Intensive versus standard physical  
17 rehabilitation therapy in the critically ill (EPICC); a multicentre, parallel-group,  
18 randomised controlled trial. *Thorax*. 2017; (5).  
19  
20  
21  
22  
23 21. Sottile PD, Nordon-Craft A, Malone D, Schenkman M, Moss M. Patient and family  
24 perceptions of physical therapy in the medical intensive care unit. *Journal of Critical*  
25 *Care*. 2015; 30, 891-895.  
26  
27  
28  
29 22. Mills, J., Bonner, A., Francis, K. (2006) The development of constructivist grounded  
30 theory. *International Journal of Qualitative Research Methods*. 5 (1).  
31  
32  
33 23. Mental Capacity Act (2005). UK legislation. Available at:  
34 <https://www.legislation.gov.uk/ukpga/2005/9/contents>  
35  
36  
37 24. Corbin, J. & Strauss, A. (2015) *Basics of qualitative research: techniques and*  
38 *procedures for developing grounded theory*. 4th edition. London  
39  
40  
41 25. Heath H, Cowley S. Developing a grounded theory approach: a comparison of Glaser  
42 and Straus. *International Journal of Nursing Studies*. 2004; 41, 141-150.  
43  
44 26. Bonde, D. (2013) *Qualitative Interviews: When enough is enough*. Research by  
45 Design.  
46  
47 27. Todres L, Galvin KT, Holloway I. The humanisation of healthcare: a value framework  
48 for qualitative research. *International Journal of Qualitative Studies on Health and*  
49 *Well-being*. 2009; 4, 68-77.  
50  
51  
52 28. Deci EL, Ryan RM. *Intrinsic motivation and self-determination in human behavior*.  
53 New York, NY: Plenum. 1985.  
54  
55  
56 29. Deci EL, Ryan RM. (2017) <http://selfdeterminationtheory.org/theory/> (Accessed:  
57 30th March 2017)  
58  
59  
60 30. Markus H & Nurius P. Possible Selves. *American Psychologist*. 1986; 41 (9), 954-969

- 1  
2  
3 31. Osman M. What are the essential cognitive requirements for prospection (thinking  
4 about the future)? *Frontiers in Psychology*. 2014;5, 626.  
5  
6  
7 32. Kean S, Salisbury LG, Rattray J, Walsh TS, Huby G, Ramsey, P. 'Intensive Care Unit  
8 (ICU) Survivorship' – a constructivist grounded theory of surviving critical illness.  
9  
10 *Journal of Clinical Nursing*. 2017. [online].  
11  
12 33. Johnston LB. Surviving Critical Illness: A Case Study in Ambiguity. *Journal of Social*  
13 *Work in End-of-Life & Palliative Care*. 2011; 7 (4), 363-382.  
14  
15 34. Darbyshire JL, Greig PR, Vollaam S, Young DJ, Hinton L. "I Can Remember Sort of Vivid  
16 People... but to Me They Were Plasticine." *Delusions on the Intensive Care Unit:*  
17 *What Do Patients Think is Going on?* *PLoS ONW*; 11 (4): e0153775.  
18  
19 35. Lindberg C, Sivberg B, Willman A, Fagerstrom C. A trajectory towards partnership in  
20 care – patients perspective of autonomy in intensive care: A qualitative study.  
21 *Intensive and Critical Care Nursing*. 2015;31: 294-302.  
22  
23 36. National Institute for Health and Clinical Excellence. Great Britain. *Rehabilitation*  
24 *After Critical Illness* Great Britain. Available at: [www.nice.org.uk](http://www.nice.org.uk). 2009.  
25  
26 37. National Institute for Health and Clinical Excellence (2017). *Rehabilitation after*  
27 *Critical illness in adults: Quality Standard*. Published 7th September 2017. Available  
28 online: [nice.org.uk/guidance/qs158](http://nice.org.uk/guidance/qs158). (Accessed: 13th September 2017)  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Topic guide****Opening**

1. Introduction
2. Consent confirmed.

**Questions**

3. Do you have any questions about the about the patient information sheet?
4. Could you tell me about the events leading up to your admission to the ICU?
5. Could you summarize, as you remember it, your stay on the ICU including the length of your stay and the procedures you experienced (e.g. surgery, tracheostomy etc)?
6. Could you describe any physical problems that you had during and after your stay, such as weakness, pain, joint stiffness etc?
7. Could you describe your rehabilitation experience?
  - Memories of rehabilitation
  - Rehabilitation equipment
  - Interaction with the therapist
  - Intensity of rehabilitation
  - Rehabilitation goal setting

**Closing**

8. Do you have any additional information you would like to add?
9. Do you have any questions?

**End****Table 1: Topic guide**

Table 2: Summary of each participant

Pseudonym	Relative present?	Age range (years)	APACHE II	Diagnosis	Length of stay, ICU (days)	Length of stay, hospital (days)	Discharge location
Richard	No	30-39	17	Acute porphyria	9	102	Long-term, inpatient rehabilitation
Martin	No	30-39	14	Drug overdose, aspiration pneumonia, rhabdomyolysis	26	32	Home, outpatient rehabilitation
Sadiq	No	50-59	22	Exacerbation of COPD	33	34	Home, full care package
Sarah	No	60-69	24	Open hernia repair- post operative MOF	115	197	Long-term, inpatient rehabilitation
Tom	No	60-69	10	Pneumonia and pulmonary embolism	10	16	Home, outpatient rehabilitation
Evan	No	60-69	15	Acute bowel obstruction- colon cancer	5	48	Home, no rehabilitation.
Sasha	Yes, daughter	50-59	10	Neuromyelitis optica	19	98	Long-term, inpatient rehabilitation
John	No	40-49	27	Influenza	33	71	Long-term, inpatient rehabilitation
George	No	50-59	12	Drug overdose- respiratory failure	25	36	Home, no rehabilitation
Michelle	No	80-89	14	Exacerbation of COPD	6	42	Declined inpatient rehabilitation- home, full care package
Jim (M)	Yes, wife	50-59	11	Food poisoning- MOF	10	18	Home, no rehabilitation
Matthew (M)	No	70-79	18	Hospital acquired pneumonia- fractured NOF	5	178	Nursing home
Caroline (F)	Yes, husband	70-79	22	Anterior resection for bowel cancer	13	63	Short-stay, inpatient rehabilitation.
Ben (M)	No	40-49	15	Drug overdose	65	107	Home, care package
David (M)	Yes, wife present	63	21	Influenza	150	232	Long-term, inpatient rehabilitation

1  
2  
3  
4  
5  
6  
7 (NOF-neck of femur; MOF-multi-organ failure; COPD-chronic obstructive pulmonary disease; APACHE II- acute  
8 physiology and chronic health evaluation II; ICU – Intensive Care Unit)  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For peer review only

**Table 3:** Supporting data

<b>Central phenomenon: Recalibration of The Self</b>	
<b>Main theme 1: "From prior self to current self"</b>	
<p><i>Sasha: "I didn't realize I couldn't walk. I thought I could and I tried to get out of bed loads of times, but up here I was weak (legs) and the top of my arms were weak as well. I couldn't do it."</i></p> <p><i>Sarah: "I don't ever look at myself in the mirror and there is a mirror in that bathroom, I just happened to catch sight of my whole body almost and I nearly died. I thought; that doesn't resemble the person that I am."</i></p>	
<b>Episodic memory loss</b>	<p><b>ICU admission</b></p> <p>Sasha: "...that's when I don't know, it's a real black after that (the emergency room)"</p> <p>Sadiq: "That is a black. That is a blank. Totally blank"</p> <p>John: "I must have been in and out of consciousness, because I don't remember anything"</p> <p>Ben: "I had a bad fall, collapsed... that's all I remember"</p> <p><b>Rehabilitation and mobilization</b></p> <p>EJC: "what was your memory of getting moving after you woke up with the tubes attached?" Ben: "I don't really have much memory of it."</p> <p>EJC: Do you remember any of the rehab on ICU?" Martin: "Not to start with, no."</p> <p>David: "It was Dan (ward physiotherapist) who taught me to sit on the edge of the bed."</p> <p>EJC: "Do you remember getting into the chair for the first time?" Michelle: "It was with Tom (the ward physical therapist)."</p>
<b>Hallucinations and delusions</b>	<p>John: "I kept thinking I could see like people with hoodies and they were like assassins, trying to get in."</p> <p>David: "I was taken into Soho (Central London) by some people and stuck under a glass floor, lying under a glass floor with formaldehyde around me. I was encased."</p> <p>Ben: "I operated on Margaret Thatchers cat and there was eight other people in the house and three of them got shot... I remember waking up with the fear that I was going to get shot."</p> <p>Carolyn: "I was trying to use my mobile (to escape), and the same number kept on pressing and I remember panicking"</p>



<p><b>Weakness</b></p>	<p>George: "Nothing, I couldn't move my hand. I couldn't move and that was really scary. Really scary."</p> <p>Richard: " I couldn't do anything. I was paralyzed from the neck down... I still felt like I had sensation in my legs and my arms, I just couldn't move them."</p> <p>Carolyn: "I couldn't even stand up. I was really very, very weak."</p> <p>John: "I couldn't do anything. I mean literally, I couldn't move, I could just barely move my fingers."</p> <p>David: "I couldn't move. I couldn't move at all. I could blink, that's about it."</p> <p>Martin: "...couldn't walk, couldn't do nothing."</p> <p>Richard: "You are reliant entirely on the people around you, for everything really... that's difficult."</p>
<p><b>Noxious cycle of ICU</b></p>	<p>Sarah: "I didn't want to do it (physiotherapy). I used to dread them coming, any excuse to get out of it. I was just so tired."</p> <p>John: "Physically tiring, emotionally... you're like 'sh*t really? I've got to do it (mobilization) now? I haven't got any energy at all."</p> <p>George: "There were some days when they'd (physiotherapist) come and they'd get me into the chair, and they'd want to do some work on the zimmer frame. They'd come back (from getting the zimmer frame) and I'd be asleep."</p> <p>Sarah: "People kept telling me to read, but I couldn't. I couldn't' actually physically read. They'd bring me the menu and I just couldn't do it, and then I'd fall asleep"</p> <p>Sarah: "...then I just accepted it (weakness), going...on the hoist and, you lose all dignity when you're in that state you just accept it, and you just let them help you as much as possible and when you've done your, you know bits of physio, exhausted, you go back to bed again, sleep again. You know it tended to be like that."</p>
<p style="text-align: center;"><b>Main theme 2: "From current self to construction of the future self"</b></p> <p>Ben: <i>"The first days when I couldn't move... I was disillusioned with the whole thing, and I thought, 'This is never going to work'... I couldn't see how anything could turn round, but I was told just to trust. But that period was very difficult because when you don't see any light at the end of the tunnel, it's difficult to sort of engage with it, and it's difficult to trust... There was plans in my head, but it's difficult to kind of have them if you think it's just a waste of time what you're doing. Now I know that there is (light at the end of the tunnel)... and I believe I'll be walking next week, they've (physiotherapists) let me believe that"</i></p>	

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

**Recovery milestones and goal setting**

Matthew: "Let the patient realize that he is not capable of doing that, or this, or whatever... don't tell him"

Tom: "Everyone's functions, and how they are improving, might not be quite so obvious to the patient."

Carolyn: "The other day the whole ward congratulated me- and even now I feel embarrassed – because I washed myself. I didn't wait until now to know how to wash myself; I thought it was so stupid."

Jim's wife: "We didn't want to set the goals, because we didn't know what goals to set"

Sadiq: "It depends on the person. If a person is shooting to the high, they might do it (achieve their goal), but sometimes shooting too much to the high might break your neck. If they are too sick, they cannot talk, you are in the dark and you have to put your own objectives."

Researcher: "What have been the things that have kept you going?" Sasha: "I think Gemma (daughter) and her dad, they've been so supportive. He's been down every day, and Gemma sometimes twice a day \*starts crying\*... sorry...I suppose if it wasn't for them, I wouldn't be... \*crying- unable to finish sentence\*

Richard: "obviously I was doing it (rehab) for me primarily, but knowing how much concern and love she has for me, and knowing how much it would mean to her and how much of a relief it would be to her... The fact that she was, you know with me for as long as she was, and as strong as she was... I don't know. I never thought my mum was that strong."

1  
2  
3 Figure legends  
4  
5

6 **Figure 1: The noxious cycle of critical illness.**  
7  
8  
9

10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For peer review only

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60




Figure 1: noxious cycle of ICU